

TRACES OF SINGULAR VALUES AND BORCHERDS PRODUCTS

CHANG HEON KIM

ABSTRACT. Let p be a prime for which the congruence group $\Gamma_0(p)^*$ is of genus zero, and j_p^* be the corresponding Hauptmodul. Let f be a nearly holomorphic modular form of weight $1/2$ on $\Gamma_0(4p)$ which satisfies some congruence condition on its Fourier coefficients. We interpret f as a vector valued modular form. Applying Borchers lifting of vector valued modular forms we construct infinite products associated to j_p^* and extend Zagier's trace formula for singular values of j_p^* . Further we investigate the twisted traces of singular values of j_p^* and construct Borchers products related to them.

1. INTRODUCTION

Let d denote a positive integer congruent to 0 or 3 modulo 4. We denote by $\mathcal{Q}_d$ the set of positive definite binary quadratic forms $Q = [a, b, c] = aX^2 + bXY + cY^2$ ($a, b, c \in \mathbb{Z}$) of discriminant $-d$, with usual action of the modular group $\Gamma = SL_2(\mathbb{Z})$. To each $Q \in \mathcal{Q}_d$, we associate its unique root $\alpha_Q \in \mathfrak{H}$ (=upper half plane). Let $j(\tau)$ ($\tau \in \mathfrak{H}$) be the elliptic modular invariant and $\mathbf{t}(d)$ be the (weighted) trace of a singular modulus of discriminant $-d$ ($= \sum_{Q \in \mathcal{Q}_d/\Gamma} \frac{1}{|\Gamma_Q|} (j(\alpha_Q) - 744)$). Here $\bar{\Gamma}_Q = \{\gamma \in \bar{\Gamma} = PSL_2(\mathbb{Z}) \mid Q \circ \gamma = Q\}$. In addition we set $\mathbf{t}(-1) = -1$, $\mathbf{t}(0) = 2$ and $\mathbf{t}(d) = 0$ for $d < -1$ or $d \equiv 1, 2 \pmod{4}$. Zagier showed the series $\sum_{d \in \mathbb{Z}} \mathbf{t}(d)q^d$ ($q = e^{2\pi i\tau}$, $\tau \in \mathfrak{H}$) is a modular form of weight $3/2$ on $\Gamma_0(4) = \{(\begin{smallmatrix} a & b \\ c & d \end{smallmatrix}) \in SL_2(\mathbb{Z}), 4|c\}$, holomorphic in $\mathfrak{H}$ and meromorphic at cusps ([18] Theorem 1).

Let $\Gamma_0(N)^*$ ($N = 1, 2, 3, \dots$) be the group generated by $\Gamma_0(N)$ and all Atkin-Lehner involutions W_e for $e|N$. Here W_e is represented by a matrix of the form $\frac{1}{\sqrt{e}} \begin{pmatrix} ex & y \\ Nz & ew \end{pmatrix}$ with

2000 *Mathematics Subject Classification*: Primary 11F03, 11F30; Secondary 11F22, 11F37, 11F50

$x, y, z, w \in \mathbb{Z}$ and $xwe - yzN/e = 1$. There are only finitely many values of N for which $\Gamma_0(N)^*$ is of genus 0. In particular, if we let $\mathfrak{S}$ denote the set of prime values for such N , then

$$\mathfrak{S} = \{2, 3, 5, 7, 11, 13, 17, 19, 23, 29, 31, 41, 47, 59, 71\}.$$

Let j_N^* be the corresponding *Hauptmodul*, whose Fourier expansion starts with $q^{-1} + 0 + a_1q + a_2q^2 + \dots$. It can be described by means of Dedekind eta function or theta functions. For example, if $N - 1$ divides 24, then the Hauptmodul j_N^* is explicitly given by

$$j_N^*(\tau) = \left(\frac{\eta(\tau)}{\eta(N\tau)} \right)^{\frac{24}{N-1}} + \frac{24}{N-1} + N^{\frac{12}{N-1}} \left(\frac{\eta(N\tau)}{\eta(\tau)} \right)^{\frac{24}{N-1}}.$$

By a result of Borcherds [1] the Fourier coefficients of j_N^* are related to certain representations of the monster simple group.

Let d be an integer ≥ 0 such that $-d$ is congruent to a square modulo $4N$. We choose an integer $\beta \pmod{2N}$ with $\beta^2 \equiv -d \pmod{4N}$ and consider the set $\mathcal{Q}_{d,N,\beta} = \{[a, b, c] \in \mathcal{Q}_d \mid a \equiv 0 \pmod{N}, b \equiv \beta \pmod{2N}\}$ on which $\Gamma_0(N)$ acts. We assume that d is not divisible as a discriminant by the square of any prime dividing N . Then the image of the root α_Q ($Q \in \mathcal{Q}_{d,N,\beta}$) in $\Gamma_0(N) \backslash \mathfrak{H}$ corresponds to Heegner point and the natural map from the quotient $\mathcal{Q}_{d,N,\beta}/\Gamma_0(N)$ to $\mathcal{Q}_d/\Gamma$ is a bijection. We define a trace $\mathbf{t}^{(N,\beta)}(d)$ by $\mathbf{t}^{(N,\beta)}(d) = \sum_{Q \in \mathcal{Q}_{d,N,\beta}/\Gamma_0(N)} \frac{1}{|\Gamma_Q|} j_N^*(\alpha_Q)$. Since j_N^* is the Hauptmodul for $\Gamma_0(N)^*$, the value of $\mathbf{t}^{(N,\beta)}(d)$ is independent of the choice of β and we can simply write $\mathbf{t}^{(N)}(d)$ instead of $\mathbf{t}^{(N,\beta)}(d)$. Zagier described $\mathbf{t}^{(N)}(d)$ as the Fourier coefficient of certain Jacobi form of weight 2 and index N for $2 \leq N \leq 6$ (see [18] Theorem 8).

Let $M_{k+1/2}^!(N)$ be the vector space consisting of *nearly holomorphic modular forms* (holomorphic in $\mathfrak{H}$ and meromorphic at cusps) of half-integral weight $k + 1/2$ on $\Gamma_0(4N)$ whose n -th Fourier coefficient vanishes unless $(-1)^k n$ is congruent to a square modulo $4N$. According to Borcherds' duality theorem (Theorem 3.1 in [4]), the only obstructions to finding such forms of weight $1/2$ are given by holomorphic vector valued modular forms

of weight $3/2$, which can be identified with elements in $J_{2,N}$ (=the space of holomorphic Jacobi forms of weight 2 and index N) (see [11] Theorem 5.1). Let N be a positive integer for which the dimension of $J_{2,N}$ is zero. Then for every integer $d \geq 0$ such that $-d$ is congruent to a square modulo $4N$, we can find a unique modular form $f_{d,N} \in M_{1/2}^!(N)$ having a Fourier expansion of the form

$$f_{d,N} = q^{-d} + \sum_{D>0} A(D, d) q^D.$$

We define a polynomial $\mathcal{H}_{d,N}(X)$ by $\prod_{Q \in \mathcal{Q}_{d,N,\beta}/\Gamma_0(N)} (X - j_N^*(\alpha_Q))^{1/|\bar{\Gamma}_Q|}$. In [13], from Zagier's trace formula ([18] Theorem 8) we derived the following product identity: for $N = 2, 3, 5, 6$

$$\mathcal{H}_{d,N}(j_N^*(\tau)) = q^{-H(d)} \prod_{u=1}^{\infty} (1 - q^u)^{A^*(u^2, d)} \quad (1)$$

where $H(d) (= \sum_{Q \in \mathcal{Q}_d/\Gamma} \frac{1}{|\bar{\Gamma}_Q|})$ is the *Hurwitz-Kronecker class number* and $A^*(D, d)$ is defined to be $2^{s(D,N)} A(D, d)$ with $s(D, N)$ the number of distinct prime factors dividing (D, N) .

Given an even lattice M in a real quadratic space of signature $(2, n)$, Borcherds lifting [3] gives a multiplicative correspondence between vector valued modular forms F of weight $1 - n/2$ with values in $\mathbb{C}[M'/M]$ (= the group ring of M'/M) and meromorphic modular forms on complex varieties $(O(2) \times O(n)) \backslash O(2, n) / \text{Aut}(M, F)$. Here $O(2, n)$ is the orthogonal group of $M \otimes \mathbb{R}$ and $\text{Aut}(M, F)$ is the subgroup of orthogonal group of M ($= \text{Aut}(M)$) leaving the form F stable under the natural action of $\text{Aut}(M)$ on M'/M . In this article we are mainly concerned with $O(2, 1)$ case. For $O(2, n)$ ($n \geq 2$) case one is referred to [5, 6]. We take M to be the direct sum of 2-dimensional unimodular Lorentzian lattice and 1-dimensional lattice generated by a vector of norm $2N$, as in Example 5.1 of [4]. If $N = 1$, each element in $M_{1/2}^!(1)$ has a natural interpretation as a vector valued modular form ([2] Lemma 14.2). Let N be a prime p for which $\Gamma_0(p)^*$ is of genus zero, i.e. $p \in \mathfrak{S}$. We will interpret a scalar valued modular form $f \in M_{1/2}^!(p)$ as a vector valued modular form

F with values in $\mathbb{C}[M'/M]$. The Borchers lifting of F is then a meromorphic modular form on $\Gamma_0(p)^*$ whose zeros or poles occur only at cusps or imaginary quadratic irrationals corresponding to Heegner points. Moreover it has an explicit form of infinite product expansion ([3] Theorem 13.3) which allows us to extend (1) to hold true for all $p \in \mathfrak{S}$. (see (16)). As a byproduct we can extend Zagier's trace formula to all primes $p \in \mathfrak{S}$ (see Corollary 3.6).

Corollary 3.6 gives a description of the coefficient of q^1 in $f_{d,N}$ as the trace of singular value of j_N^* . In [18] §7 when the level N is equal to 1 and $D > 1$, Zagier described the D -th coefficient of $f_{d,1}$ in terms of the relative trace of singular modulus of discriminant $-dD$ from the Hilbert class field of $\mathbb{Q}(\sqrt{-dD})$ to its real quadratic subfield $\mathbb{Q}(\sqrt{D})$. In §4 we will investigate the analogue of this in higher level cases and construct Borchers products (see Theorem 4.4).

2. PRELIMINARIES

2.1. Modular forms of half-integral weight. Here we recall some basic definitions. Let $\mathfrak{G}$ be the group consisting of all pairs $(A, \psi(\tau))$, where $A = \begin{pmatrix} a & b \\ c & d \end{pmatrix} \in GL_2^+(\mathbb{R})$ and $\psi(\tau)$ is a complex valued function holomorphic on $\mathfrak{H}$ satisfying $|\psi(\tau)| = (\det A)^{-1/4} |c\tau + d|^{1/2}$, with group law defined by $(A, \psi_1(\tau))(B, \psi_2(\tau)) = (AB, \psi_1(B\tau)\psi_2(\tau))$. If $f : \mathfrak{H} \rightarrow \mathbb{C}$ and $\xi = (A, \psi(\tau)) \in \mathfrak{G}$, we put $f|_{[\xi]_{k+1/2}} = f|_{\xi} = \psi(\tau)^{-2k-1} f(A\tau)$. Then from definition it follows that $f|_{\xi_1}|_{\xi_2} = f|_{\xi_1\xi_2}$. There is a monomorphism $\Gamma_0(4) \rightarrow \mathfrak{G}$ given by $A \mapsto A^* := (A, j(A, \tau))$, where $j(A, \tau) = \left(\frac{c}{d}\right) \left(\frac{-1}{d}\right)^{-1/2} (c\tau + d)^{1/2}$ if $A = \begin{pmatrix} a & b \\ c & d \end{pmatrix}$. A complex valued holomorphic function f on $\mathfrak{H}$ is said to be a *modular form of half-integral weight $k + 1/2$ on $\Gamma_0(4N)$* if it satisfies $f|_{A^*} = f$ for every $A \in \Gamma_0(4N)$, and is holomorphic at the cusps.

2.2. Jacobi forms. A (*holomorphic*) *Jacobi form of weight k and index N* is defined to be a holomorphic function $\phi : \mathfrak{H} \times \mathbb{C} \rightarrow \mathbb{C}$ satisfying the two transformation laws

$$\begin{aligned} \phi\left(\frac{a\tau + b}{c\tau + d}, \frac{z}{c\tau + d}\right) &= (c\tau + d)^k e^{2\pi i N \frac{cz^2}{c\tau + d}} \phi(\tau, z) \quad (\forall \begin{pmatrix} a & b \\ c & d \end{pmatrix} \in SL_2(\mathbb{Z})), \\ \phi(\tau, z + \lambda\tau + \mu) &= e^{-2\pi i N(\lambda^2\tau + 2\lambda z)} \phi(\tau, z) \quad (\forall \begin{pmatrix} \lambda & \mu \end{pmatrix} \in \mathbb{Z}^2) \end{aligned}$$

and having a Fourier expansion of the form

$$\phi(\tau, z) = \sum_{\substack{n, r \in \mathbb{Z} \\ 4Nn - r^2 \geq 0}} c(n, r) q^n \zeta^r \quad (q = e^{2\pi i \tau}, \zeta = e^{2\pi i z}), \quad (2)$$

where the coefficient $c(n, r)$ depends only on $4Nn - r^2$ and on the residue class of $r \pmod{2N}$ ([11] Theorem 2.2). In (2), if the condition $4Nn - r^2 \geq 0$ is deleted, we obtain a *nearly holomorphic Jacobi form*.

Let $J_{k,N}^!$ be the space of nearly holomorphic Jacobi forms of weight k and index N . Let $J_{*,*}^!$ be the ring of all nearly holomorphic Jacobi forms and $J_{ev,*}^!$ its even weight subring. Then $J_{ev,*}^!$ is the free polynomial algebra over $M_*^!(\Gamma) = \mathbb{C}[E_4, E_6, \Delta^{-1}]/(E_4^3 - E_6^2 = 1728\Delta)$ on two generators $a = \tilde{\phi}_{-2,1}(\tau, z) \in J_{-2,1}^!$ and $b = \tilde{\phi}_{0,1}(\tau, z) \in J_{0,1}^!$ (see [11] §9). Let $N \in \{1\} \cup \mathfrak{S}$ and $k = 2$. There are Jacobi forms $\phi_{D,N} \in J_{2,N}^!$ uniquely characterized by the requirement that they have Fourier coefficients $c(n, r) = B(D, 4Nn - r^2)$ which depend only on the discriminant $r^2 - 4Nn$, with $B(D, -D) = 1$, $B(D, d) = 0$ if $d = 4Nn - r^2 < 0, \neq -D$ and $B(D, 0) = -2$ or 0 according as D is a square or non-square. The uniqueness of $\phi_{D,N}$ is obvious since the difference of any two functions satisfying the definition of $\phi_{D,N}$ would be an element of $J_{2,N}$, which is of dimension zero (see [11] p.118). For the existence the structure theorem enables us to express $\phi_{D,N}$ as a linear combination of $a^i b^{N-i}$ ($i = 0, \dots, N$) over $M_*^!(\Gamma)$.

Through the article we adopt the following notations:

- β : an element in $\mathbb{Z}/2N\mathbb{Z}$.

- $\bar{\Gamma}'$: the image of Γ' in $PSL_2(\mathbb{R})$.
- $\bar{\Gamma}'_Q := \{\gamma \in \bar{\Gamma}' \mid Q \circ \gamma = Q\}$.
- When we write $\mathcal{Q}_{d,N,\beta}$, β is always assumed to satisfy $\beta^2 \equiv -d \pmod{4N}$.
- $e(x) := \exp(2\pi i x)$.
- $\zeta_n := e(1/n) = \exp(2\pi i/n)$.
- $s(D, N)$: the number of prime factors dividing (D, N) .
- U_m : the Hecke operator defined by $\sum_{n \in \mathbb{Z}} c(n)q^n \mid_{U_m} = \sum_{n \in \mathbb{Z}} c(mn)q^n$.
- $E_k(\tau)$: the normalized Eisenstein series of weight k , equal to $1 - (2k/B_k) \sum_{n>0} \sigma_{k-1}(n)q^n$ where B_k is the k -th Bernoulli number defined by $\sum_{k \in \mathbb{Z}} B_k t^k/k! = t/(e^t - 1)$ and $\sigma_{k-1}(n) = \sum_{d|n, d>0} d^{k-1}$.

3. VECTOR VALUED MODULAR FORMS

Definition 3.1. Let M be an even lattice of signature $(2, n)$ equipped with a non-degenerate quadratic form $q(x) = \frac{1}{2}(x, x)$. Let M' be the dual lattice of M and $\mathbb{C}[M'/M]$ be the group ring of M'/M with basis $\mathbf{e}_\gamma$ for each $\gamma \in M'/M$. We define a *vector valued modular form of weight k and of type ρ_M* to be a holomorphic function $F(\tau) = \sum_{\gamma \in M'/M} h_\gamma(\tau) \mathbf{e}_\gamma$ on the upper half plane $\mathfrak{H}$ with values in $\mathbb{C}[M'/M]$ such that

$$\begin{aligned} h_\gamma(\tau + 1) &= e((\gamma, \gamma)/2) h_\gamma(\tau) \\ h_\gamma(-1/\tau) &= \frac{\sqrt{i}^{n-2}}{\sqrt{|M'/M|}} \sqrt{\tau}^k \sum_{\delta \in M'/M} e(-(\gamma, \delta)) h_\delta(\tau) \\ h_{-\gamma}(\tau) &= (-1)^k i^{n-2} h_\gamma(\tau). \end{aligned}$$

For a fixed positive integer N , we take M to be the 3 dimensional even lattice of all symmetric matrices $v = \begin{pmatrix} A & B \\ B & C \end{pmatrix}$ with $NA, B, C \in \mathbb{Z}$ with the norm $(v, v) = -2N \det(v)$. The lattice M splits as the direct sum of the 2 dimensional hyperbolic unimodular even lattice $II_{1,1} \approx \mathbb{Z} \begin{pmatrix} 1/N & 0 \\ 0 & 0 \end{pmatrix} + \mathbb{Z} \begin{pmatrix} 0 & 0 \\ 0 & 1 \end{pmatrix}$ and a lattice generated by $\begin{pmatrix} 0 & 1 \\ 1 & 0 \end{pmatrix}$ of norm $2N$. Thus

the dual lattice M' is the set of matrices $\begin{pmatrix} A & B \\ B & C \end{pmatrix}$ with $NA, 2NB, C \in \mathbb{Z}$, and M'/M can be identified with $\mathbb{Z}/2N\mathbb{Z}$ by mapping a matrix of M' to the value of $B \in \frac{1}{2N}\mathbb{Z}/\mathbb{Z} \approx \mathbb{Z}/2N\mathbb{Z}$. The group $\Gamma_0(N)^*$ acts on the lattice M as an orthogonal transformation by $v \mapsto XvX^t$ for $v \in M$ and $X \in \Gamma_0(N)^*$. Moreover it is not difficult to verify that the special orthogonal group of M is $\Gamma_0(N)^*$.

In the following we will interpret a scalar valued modular form in $M_{1/2}^!(N)$ as a vector valued modular form of weight $1/2$ and type ρ_M . For simplicity we will restrict ourselves to the case N is a prime p in the set $\mathfrak{S}$. Let $f = \sum_{n \in \mathbb{Z}} c(n)q^n \in M_{1/2}^!(p)$. For $\beta \in \mathbb{Z}/2p\mathbb{Z}$, we set

$$h_\beta(\tau) = 2^{s(\beta,p)-1} \sum_{n \equiv \beta^2 (4p)} c(n)q^{n/4p}$$

where $s(\beta, p) = 1$ if $\beta \equiv 0, p \pmod{2p}$ and 0 otherwise. Then clearly we have

$$h_\beta(\tau + 1) = \zeta_{4p}^{\beta^2} h_\beta(\tau). \quad (3)$$

For j prime to $4p$, we consider $f_j = f|_{\left(\begin{pmatrix} 1 & j \\ 0 & 4p \end{pmatrix}, (4p)^{1/4}\right)}|_{W_{4p}}$ where $W_{4p} = \left(\begin{pmatrix} 0 & -1 \\ 4p & 0 \end{pmatrix}, (4p)^{1/4}\sqrt{-i\tau}\right)$.

By definition $f_j = f|_{\left(\begin{pmatrix} 4pj & -1 \\ 16p^2 & 0 \end{pmatrix}, 2\sqrt{-pi\tau}\right)}$. Let b and d be integers satisfying $jd - 4pb = 1$ and

$$\psi_j = \left(\frac{4p}{j}\right) \sqrt{\left(\frac{-1}{j}\right)} \zeta_8^{-1}. \text{ Then it holds that } \left(\begin{pmatrix} j & b \\ 4p & d \end{pmatrix}\right)^* \left(\begin{pmatrix} 4p & -d \\ 0 & 4p \end{pmatrix}, \psi_j\right) = \left(\begin{pmatrix} 4pj & -1 \\ 16p^2 & 0 \end{pmatrix}, 2\sqrt{-pi\tau}\right).$$

Thus $f_j = f|_{\left(\begin{pmatrix} j & b \\ 4p & d \end{pmatrix}\right)^* \left(\begin{pmatrix} 4p & -d \\ 0 & 4p \end{pmatrix}, \psi_j\right)} = \psi_j^{-1} f\left(\tau - \frac{j^{-1}}{4p}\right)$, from which it follows that

$$f|_{\left(\begin{pmatrix} 1 & j \\ 0 & 4p \end{pmatrix}, (4p)^{1/4}\right)}|_{W_{4p}} = \psi_j^{-1} f\left(\tau - \frac{j^{-1}}{4p}\right) \quad (4)$$

where j^{-1} denotes the inverse of j in $(\mathbb{Z}/4p\mathbb{Z})^*$. The left hand side of (4) is

$$\begin{aligned} (4p)^{-1/4} f\left(\frac{\tau + j}{4p}\right)|_{W_{4p}} &= (4p)^{-1/4} \left(\sum_{n \in \mathbb{Z}} c(n) \zeta_{4p}^{nj} q^{n/4p}\right)|_{W_{4p}} = (4p)^{-1/4} \left(\sum_{\beta(2p)} \zeta_{4p}^{\beta^2 j} h_\beta\right)|_{W_{4p}} \\ &= (4p)^{-1/2} (\sqrt{-i\tau})^{-1} \left(\sum_{\beta(2p)} \zeta_{4p}^{\beta^2 j} h_\beta\right) \left(-\frac{1}{4p\tau}\right). \end{aligned} \quad (5)$$

On the other hand, the right hand side of (4) is

$$\psi_j^{-1} \left(\sum_{n \in \mathbb{Z}} c(n) \zeta_{4p}^{-nj^{-1}} q^n \right) = \psi_j^{-1} \left(\sum_{\beta(2p)} \zeta_{4p}^{-\beta^2 j^{-1}} h_\beta \right) (4p\tau). \quad (6)$$

Replacing τ by $\tau/4p$ in (5), (6) and recalling the definition of ψ_j we obtain the following identity:

$$\sum_{\beta(2p)} \zeta_{4p}^{\beta^2 j} h_\beta(-1/\tau) = \left(\frac{4p}{j} \right) \sqrt{\left(\frac{-1}{j} \right)}^{-1} \sqrt{\tau} \cdot \sum_{\beta(2p)} \zeta_{4p}^{-\beta^2 j^{-1}} h_\beta(\tau). \quad (7)$$

Let R be a $2p \times 2p$ matrix defined by $R = \frac{\zeta_8^{-1}}{\sqrt{2p}} \left(\zeta_{2p}^{-l\beta} \right)_{0 \leq l, \beta < 2p}$. To show that $\sum_{\beta(2p)} h_\beta \mathbf{e}_\beta$ is a vector valued modular form, it is necessary to check that

$$\begin{pmatrix} h_0 \\ \vdots \\ h_\beta \\ \vdots \end{pmatrix} (-1/\tau) = \sqrt{\tau} R \begin{pmatrix} h_0 \\ \vdots \\ h_\beta \\ \vdots \end{pmatrix} (\tau). \quad (8)$$

Since $h_\beta = h_{-\beta}$, the above identity is equivalent to

$$A \begin{pmatrix} h_0 \\ \vdots \\ h_\beta \\ \vdots \end{pmatrix} (-1/\tau) = \sqrt{\tau} AR \begin{pmatrix} h_0 \\ \vdots \\ h_\beta \\ \vdots \end{pmatrix} (\tau) \quad (9)$$

for some matrix A with $2p$ columns of which the first $p+1$ ones are linearly independent and span the other ones. We take $A = \left(\zeta_{4p}^{\beta^2 j_l} \right)_{\substack{1 \leq l \leq \varphi(4p) \\ 0 \leq \beta < 2p}}$ where j_l is the l -th largest element in the set $\{j \mid 1 \leq j \leq 4p \text{ and } (j, 4p) = 1\}$. We can check that the rank of A is $p+1$ unless $p = 2$. Moreover from the Gauss quadratic sum identity ([15] Chapter IV, Section 3) it follows that $AR = \left(\left(\frac{4p}{j_l} \right) \sqrt{\left(\frac{-1}{j_l} \right)}^{-1} \zeta_{4p}^{-\beta^2 j_l^{-1}} \right)_{\substack{1 \leq l \leq \varphi(4p) \\ 0 \leq \beta < 2p}}$. Now the identity (9) follows from (7) except for the case $p = 2$.

Remark 3.2. When $p = 2$, the matrix A is $\begin{pmatrix} 1 & \zeta_8 & -1 & \zeta_8 \\ 1 & \zeta_8^3 & -1 & \zeta_8^3 \\ 1 & \zeta_8^5 & -1 & \zeta_8^5 \\ 1 & \zeta_8^7 & -1 & \zeta_8^7 \end{pmatrix}$ and its rank is 2. If we take $j = 1, 5$ in (7), we obtain the following two identities:

$$h_0(-1/\tau) + 2\zeta_8 h_1(-1/\tau) - h_2(-1/\tau) = \sqrt{\tau}(h_0(\tau) + 2\zeta_8^{-1}h_1(\tau) - h_2(\tau)) \quad (10)$$

$$h_0(-1/\tau) - 2\zeta_8 h_1(-1/\tau) - h_2(-1/\tau) = -\sqrt{\tau}(h_0(\tau) - 2\zeta_8^{-1}h_1(\tau) - h_2(\tau)). \quad (11)$$

Thus in order to determine $h_\beta(-1/\tau)$ in terms of $h_\gamma(\tau)$'s we need one more identity. We consider

$$\begin{aligned} 2(f|_{U_2})|_{((\begin{smallmatrix} 1 & 0 \\ 4 & 1 \end{smallmatrix}), \sqrt{4\tau+1})} &= \sum_{j=0}^1 \sqrt[4]{2} f|_{((\begin{smallmatrix} 1 & j \\ 0 & 2 \end{smallmatrix}), \sqrt[4]{2})((\begin{smallmatrix} 1 & 0 \\ 4 & 1 \end{smallmatrix}), \sqrt{4\tau+1})} = \sum_{j=0}^1 \sqrt[4]{2} f|_{((\begin{smallmatrix} 1+4j & j \\ 8 & 2 \end{smallmatrix}), \sqrt[4]{2}\sqrt{4\tau+1})} \\ &= \sqrt[4]{2} f|_{(\begin{smallmatrix} 1 & 0 \\ 8 & 1 \end{smallmatrix})^*(\begin{smallmatrix} 1 & 0 \\ 0 & 2 \end{smallmatrix}), \sqrt[4]{2})} + \sqrt[4]{2} f|_{(\begin{smallmatrix} 5 & 3 \\ 8 & 5 \end{smallmatrix})^*(\begin{smallmatrix} 1 & -1 \\ 0 & 2 \end{smallmatrix}), \sqrt[4]{2})} \\ &= f\left(\frac{\tau}{2}\right) - f\left(\frac{\tau+1}{2}\right). \end{aligned}$$

In terms of $S = ((\begin{smallmatrix} 0 & -1 \\ 1 & 0 \end{smallmatrix}), \sqrt{\tau})$ and $T = ((\begin{smallmatrix} 1 & 1 \\ 0 & 1 \end{smallmatrix}), 1)$ we can rewrite the above as $2(f|_{U_2})|_{ST^{-4}S^{-1}} = f\left(\frac{\tau}{2}\right) - f\left(\frac{\tau+1}{2}\right)$. This yields

$$(f|_{U_2})|_S = \frac{1}{2} \left(f\left(\frac{\tau}{2}\right) - f\left(\frac{\tau+1}{2}\right) \right) |_{ST^4}. \quad (12)$$

The left hand side of (12) can be written as

$$(h_0(4\tau) + h_2(4\tau))|_S = (h_0(-4/\tau) + h_2(-4/\tau))\sqrt{\tau}^{-1}.$$

Meanwhile the right hand side of (12) turns out to be

$$\begin{aligned} 2h_1(4\tau)|_{ST^4} &= 2(h_1(-4/\tau)\sqrt{\tau}^{-1})|_{T^4} \\ &= 2 \left(\frac{\zeta_8^{-1}}{2} \sqrt{\tau/4} (h_0(\tau/4) - h_2(\tau/4)) \sqrt{\tau}^{-1} \right) |_{T^4} \text{ by (10) and (11)} \\ &= \frac{\zeta_8^{-1}}{2} (h_0(\tau/4) + h_2(\tau/4)) \text{ by (3)}. \end{aligned}$$

Replacing τ by 4τ we come up with

$$h_0(-1/\tau) + h_2(-1/\tau) = \zeta_8^{-1} \sqrt{\tau} (h_0(\tau) + h_2(\tau)). \quad (13)$$

Now from (10), (11) and (13) we can verify (8) in the case $p = 2$, too.

Now we will work out Borcherds lifting in $O(2, 1)$ case. With M as before, we let $G(M)$ denote the *Grassmannian* of M , i.e. the space of all two dimensional positive definite subspaces of $M \otimes \mathbb{R}$ acted on by the orthogonal group $O(2, 1)$. In our case the Grassmannian $G(M)$ can be identified with the upper half plane by mapping $\tau \in \mathfrak{H}$ to the two dimensional positive definite space spanned by the real and imaginary parts of the norm zero vector $\begin{pmatrix} \tau^2 & \tau \\ \tau & 1 \end{pmatrix}$. Let $F = \sum_{\beta} h_{\beta} \mathbf{e}_{\beta}$ be a vector valued modular form associated to $f = \sum_{n \in \mathbb{Z}} c(n) q^n \in M_{1/2}^!(p)$. We write $\sum_{n \in \mathbb{Z}} c_{\beta}(n/4p) e(n\tau/4p)$ for the Fourier expansion of $h_{\beta}(\tau)$. Theorem 13.3 of [3] implies that there is a meromorphic function Ψ_f on $G(M)$ that have infinite product expansion (explicitly given by Item 5 in Theorem 13.3 of [3]) with the following properties:

1. Ψ_f is an automorphic form of weight $c_0(0)/2$ for some unitary character of $\text{Aut}(M, F)$.
2. The only zeros or poles of Ψ_f lie on the divisors $\lambda^{\perp}$ for $\lambda \in M$, $\lambda^2 < 0$ and are zeros of order $\sum_{\substack{0 < x \in \mathbb{R} \\ x\lambda \in M'}} c_{x\lambda}(x^2 \lambda^2 / 2)$.

We can translate the above in terms of usual modular forms as follows:

Theorem 3.3. *With the same notation as above, we have,*

(i) *for some h ,*

$$\Psi_f(\tau) = q^{-h} \prod_{n>0} (1 - q^n)^{c_{\beta}(n^2/4p)}, \quad (\text{here for each } n > 0, \beta \equiv n \pmod{2p})$$

is a meromorphic modular form of weight $c_0(0)$ for some character of $\Gamma_0(p)^$.*

(ii) *The possible zeros or poles of Ψ_f occur either at cusps or at imaginary quadratic irrationals $\alpha \in \mathfrak{H}$ satisfying $pA\alpha^2 + B\alpha + C = 0$ with $(A, B, C) = 1$. Let $\delta = B^2 - 4pAC < 0$.*

Then the multiplicity of the zero of Ψ_f at α is given by $\sum_{n>0} c_\beta(\delta n^2/4p)$ (here for each $n > 0$, β is chosen such that $\beta^2 \equiv \delta n^2 \pmod{4p}$).

Recall the nearly holomorphic modular form $f_{d,p}$ in $M_{1/2}^!(p)$ with Fourier expansion of the form $q^{-d} + \sum_{D \geq 1} A(D, d)q^D$. We write $2f_{d,p} = \sum_{n \in \mathbb{Z}} c(n)q^n$. By Theorem 3.3 (i),

$$\Psi_{2f_{d,p}} = q^{-h} \prod_{u=1}^{\infty} (1 - q^u)^{A^*(u^2, d)} \quad (\text{for some } h) \quad (14)$$

is a meromorphic modular form of weight 0 for some character of $\Gamma_0(p)^*$. Choose an integer $\beta \pmod{2p}$ with $\beta^2 \equiv -d \pmod{4p}$. From Theorem 3.3 (ii) it follows that the zeros of $\Psi_{2f_{d,p}}$ occur at α_Q for each $Q \in \mathcal{Q}_{d,p,\beta}$ with multiplicity

$$c_\beta(-d/4p) = 2^{s(\beta,p)-1} c(-d) = \begin{cases} 1 & \Leftrightarrow p \nmid \beta \Leftrightarrow \beta \not\equiv -\beta \pmod{2p} \Leftrightarrow Q \circ W_p \notin \mathcal{Q}_{d,p,\beta} \\ 2 & \Leftrightarrow p \mid \beta \Leftrightarrow \beta \equiv -\beta \pmod{2p} \Leftrightarrow Q \circ W_p \in \mathcal{Q}_{d,p,\beta}. \end{cases}$$

On the other hand, the multiplicity of the zero of $j_p^*(\tau) - j_p^*(\alpha_Q)$ at α_Q is

$$|\bar{\Gamma}_0(p)_Q^*| = \begin{cases} |\bar{\Gamma}_0(p)_Q|, & \text{if } Q \circ W_p \neq Q \pmod{\Gamma_0(p)}; \\ 2|\bar{\Gamma}_0(p)_Q|, & \text{if } Q \circ W_p = Q \pmod{\Gamma_0(p)}. \end{cases}$$

From this observation we come up with

$$\Psi_{2f_{d,p}} = \prod_{Q \in \mathcal{Q}_{d,p,\beta}/\Gamma_0(p)} (j_p^*(\tau) - j_p^*(\alpha_Q))^{\frac{1}{|\bar{\Gamma}_0(p)_Q|}} \quad (15)$$

Comparing (14) with (15) we get

$$q^{-H_{p,\beta}(d)} \prod_{u=1}^{\infty} (1 - q^u)^{A^*(u^2, d)} = \prod_{Q \in \mathcal{Q}_{d,p,\beta}/\Gamma_0(p)} (j_p^*(\tau) - j_p^*(\alpha_Q))^{\frac{1}{|\bar{\Gamma}_0(p)_Q|}} \quad (16)$$

where $H_{p,\beta}(d) = \sum_{Q \in \mathcal{Q}_{d,p,\beta}/\Gamma_0(p)} \frac{1}{|\bar{\Gamma}_0(p)_Q|}$. Note that if d is not divisible as a discriminant by p^2 , then $|\mathcal{Q}_{d,p,\beta}/\Gamma_0(p)| = |\mathcal{Q}_d/\Gamma|$ and $|\bar{\Gamma}_0(p)_Q| = |\bar{\Gamma}_Q|$. Therefore in this case $H_{p,\beta}(d)$ is none other than $H(d)$.

Example 3.4. Let $p = 2, d = 16$ and $\beta \equiv 0 \pmod{4}$. A set of representatives for $\mathcal{Q}_{16,2,0}/\Gamma_0(2)$ is given by $Q_1 = [4, -4, 2]$, $Q_2 = [2, 0, 2]$ and $Q_3 = [4, 0, 1]$. We observe that $|\bar{\Gamma}_0(2)_{Q_i}| = 2$ if $i = 1$ and 1 otherwise. Up to modulo $\Gamma_0(2)$, the involution W_2 fixes Q_1 and interchanges Q_2 with Q_3 . Thus, by (15), $\Psi_{2f_{16,2}} = (j_2^*(\tau) - j_2^*(\alpha_{Q_1}))^{1/2}(j_2^*(\tau) - j_2^*(\alpha_{Q_2}))^2$.

Let $f = \sum_{D \in \mathbb{Z}} A(D)q^D \in M_{1/2}^!(p)$. As before, we set $h_\beta(\tau) = 2^{s(\beta,p)-1} \sum_{D \equiv \beta^2(4p)} A(D)q^{D/4p}$ for each $\beta \in \mathbb{Z}/2p\mathbb{Z}$. By (8)

$$h_\beta(-1/\tau) = \frac{\sqrt{i}^{-1}\sqrt{\tau}}{\sqrt{2p}} \sum_{j \pmod{2p}} e(-\beta j/2p) h_j(\tau).$$

Let $\phi \in J_{2,p}^!$. The Fourier coefficient $c(n, r)$ of $q^n \zeta^r$ in ϕ depends only on the discriminant $r^2 - 4pn$ and therefore we can write $c(n, r) = B(4pn - r^2)$. For $\mu \in \mathbb{Z}/2p\mathbb{Z}$, we set $\tilde{h}_\mu = \sum_{d \equiv -\mu^2(4p)} B(d)q^{d/4p}$. The formula (9) in [11] §5 says

$$\tilde{h}_\mu(-1/\tau) = \frac{\sqrt{i} \sqrt[3]{\tau}}{\sqrt{2p}} \sum_{k \pmod{2p}} e(\mu k/2p) \tilde{h}_k(\tau).$$

Let $g = \sum_{d \in \mathbb{Z}} B(d)q^d$ with $B(d)$ defined to be zero unless $-d$ is a square modulo $4p$. From definition it holds that $f = \sum_{\beta \pmod{2p}} h_\beta(4p\tau)$ and $g = \sum_{\mu \pmod{2p}} 2^{s(\mu,p)-1} \tilde{h}_\mu(4p\tau)$. Moreover $(fg)|_{U_{4p}} = \sum_{i \pmod{2p}} (h_i \tilde{h}_i)(\tau)$. Thus $(fg)|_{U_{4p}}(\tau + 1) = (fg)|_{U_{4p}}(\tau)$ and

$$\begin{aligned} (fg)|_{U_{4p}}(-1/\tau) &= \sum_{i \pmod{2p}} (h_i \tilde{h}_i)(-1/\tau) = \frac{\tau^2}{2p} \sum_{i,j,k \pmod{2p}} e(-ij/2p) e(ik/2p) (h_j \tilde{h}_k)(\tau) \\ &= \frac{\tau^2}{2p} \sum_{j,k \pmod{2p}} \sum_{i \pmod{2p}} e(i(-j+k)/2p) (h_j \tilde{h}_k)(\tau) \\ &= \tau^2 \sum_{j \pmod{2p}} (h_j \tilde{h}_j)(\tau) \quad \text{since} \quad \sum_{i \pmod{2p}} e(i(-j+k)/2p) = \begin{cases} 2p, & \text{if } j \equiv k \pmod{2p} \\ 0, & \text{otherwise} \end{cases} \\ &= \tau^2 (fg)|_{U_{4p}}. \end{aligned}$$

$(fg)|_{U_{4p}}$ is then invariant under the action of $SL_2(\mathbb{Z})$ with a pole only at ∞ and of weight 2. Thus $(fg)|_{U_{4p}}$ can be written as the derivative of some polynomial in j . In particular the constant term of the q -expansion of fg vanishes. Applying this to $f = f_{d,p}$ and $\phi = \phi_{D,p}$ gives rise to the following lemma:

Lemma 3.5. *Let $d = 4pn - r^2$ for some integers n and r . Let $A(D, d)$ (resp. $B(D, d)$) be the coefficient of q^D (resp. $q^n \zeta^r$) in $f_{d,p} \in M_{1/2}^!(p)$ (resp. $\phi_{D,p} \in J_{2,p}^!$). Then $A(D, d) = -B(D, d)$.*

Corollary 3.6. *Let p be a prime for which $\Gamma_0(p)^*$ is of genus zero. For each natural number d which is congruent to a square modulo $4p$, let $\mathbf{t}^{(p)}(d) = \sum_{Q \in \mathcal{Q}_{d,p,\beta}/\Gamma_0(p)} \frac{1}{|\Gamma_0(p)_Q|} j_p^*(\alpha_Q)$. We also put $\mathbf{t}^{(p)}(-1) = -1$, $\mathbf{t}^{(p)}(0) = 2$ and $\mathbf{t}^{(p)}(d) = 0$ for $d < -1$. Then the series $\sum_{n,r} \mathbf{t}^{(p)}(4pn - r^2) q^n \zeta^r$ becomes a nearly holomorphic Jacobi form of weight 2 and index p .*

Proof. In (16) if we compare the q -expansions on both sides, then we obtain $\mathbf{t}^{(p)}(d) = A^*(1, d) = A(1, d)$ which is equal to $-B(1, d)$ by Lemma 3.5. Now the series $\sum_{n,r} \mathbf{t}^{(p)}(4pn - r^2) q^n \zeta^r$ coincides with $-\phi_{1,p}$. $\square$

4. TWISTED TRACES OF SINGULAR VALUES

Let D and $-d$ be a positive and a negative discriminant, respectively. For simplicity we suppose them to be fundamental and coprime. Let $K = \mathbb{Q}(\sqrt{-Dd})$. Let $p_1, \dots, p_r$ be odd primes dividing Dd and put $p_i^* = (-1)^{(p_i-1)/2} p_i$ for $i = 1, \dots, r$. By class field theory the genus field M of K is expressed as $K(\sqrt{p_1^*}, \dots, \sqrt{p_r^*})$ and is contained in the Hilbert class field H of K (see [10] Theorem 6.1). We have obvious inclusion $N = K(\sqrt{D}) \subset M \subset H$. The genus character $\chi = \chi_{D,-d}$ assigns to any quadratic form Q of discriminant $-dD$ a value ± 1 defined by $\chi(Q) = \left(\frac{D}{q}\right) = \left(\frac{-d}{q}\right)$ where q is any prime represented by Q and not dividing Dd . This is independent of the choice of q . We further assume D and $-d$ to be

congruent to a square modulo $4p$. We define the “twisted trace” $\mathbf{t}^{(p)}(D, d)$ by

$$\mathbf{t}^{(p)}(D, d) = \sum_{Q \in \mathcal{Q}_{dD, p, \beta} / \Gamma_0(p)} \chi(Q) j_p^*(\alpha_Q).$$

Proposition 4.1. (i) *The definition of $\mathbf{t}^{(p)}(D, d)$ is independent of the choice of β .*

(ii) *Let p be a prime such that $p - 1$ divides 24. Then $\frac{1}{\sqrt{D}} \mathbf{t}^{(p)}(D, d)$ is a rational integer.*

Proof. (i) Since W_p induces a bijection between $\mathcal{Q}_{dD, p, \beta} / \Gamma_0(p)$ and $\mathcal{Q}_{dD, p, -\beta} / \Gamma_0(p)$, it is enough to show that $\chi(Q \circ W_p) = \chi(Q)$ and $j_p^*(\alpha_{Q \circ W_p}) = j_p^*(\alpha_Q)$ for each $Q \in \mathcal{Q}_{dD, p, \beta}$. The latter is clear since j_p^* is the Hauptmodul for $\Gamma_0(p)^*$. The invariance of χ under W_p follows from Proposition 1 in p.508 of [12].

(ii) It is well-known that $j(\alpha_Q)$ is an algebraic integer. By [7] Remark 1.5.3 and Appendix 1 there is a modular relation between j and j_p^* which allows j_p^* to be integral over $\mathbb{Z}[j]$. Therefore $j_p^*(\alpha_Q)$ is an algebraic integer, too. Let $j_{0,p}$ be the Hauptmodul for $\Gamma_0(p)$. Then j_p^* can be written as $j_{0,p}(\tau) + j_{0,p}(W_p \tau)$ up to some constant ($\in \mathbb{Z}$). By [7] Theorem 3.7.5, $j_{0,p}(\alpha_Q)$ (resp. $j_{0,p}(W_p \alpha_Q)$) generates Hilbert class field H of K so that $j_p^*(\alpha_Q)$ belongs to H . The set of forms Q with $\chi(Q) = +1$ is identified with $\text{Gal}(H/N)$. Therefore the sum $\sum_{\chi(Q)=+1} j_p^*(\alpha_Q)$ is the trace of the algebraic integer $j_p^*(\alpha_Q)$ from H to N . Since this number is invariant under the complex conjugation, it belongs to $\mathbb{Q}(\sqrt{D})$. If we write this number as $\lambda = \frac{1}{2}(a + b\sqrt{D})$ with $a, b \in \mathbb{Z}$, then the sum over all $j_p^*(\alpha_Q)$ is equal to $\text{Tr}_{H/K}(j_p^*(\alpha_Q)) = \text{Tr}_{N/K} \circ \text{Tr}_{H/N}(j_p^*(\alpha_Q)) = \text{Tr}_{N/K}(\lambda) = a$. Hence the sum $\mathbf{t}^{(p)}(D, d)$ is equal to $-a + 2\lambda = b\sqrt{D}$. $\square$

Here are some numerical examples with $p = 2$ and $p = 3$:

$$\frac{1}{\sqrt{17}} \mathbf{t}^{(2)}(17, 4) = \frac{1}{\sqrt{17}} (j_2^*(\alpha_{[18, 2, 1]}) + j_2^*(\alpha_{[2, 2, 9]}) - j_2^*(\alpha_{[6, -2, 3]}) - j_2^*(\alpha_{[6, 2, 3]})) = -204800,$$

$$\frac{1}{\sqrt{8}} \mathbf{t}^{(2)}(8, 7) = \frac{1}{\sqrt{8}} (j_2^*(\alpha_{[14, 0, 1]}) + j_2^*(\alpha_{[2, 0, 7]}) - j_2^*(\alpha_{[6, 4, 3]}) - j_2^*(\alpha_{[10, 8, 3]})) = 90112,$$

and

$$\frac{1}{\sqrt{13}} \mathbf{t}^{(3)}(13, 3) = \frac{1}{\sqrt{13}} (j_3^*(\alpha_{[12, 3, 1]}) + j_3^*(\alpha_{[3, 3, 4]}) - j_3^*(\alpha_{[6, 3, 2]}) - j_3^*(\alpha_{[15, 9, 2]})) = -378,$$

$$\frac{1}{\sqrt{13}}\mathbf{t}^{(3)}(13, 8) = \frac{1}{\sqrt{13}}(j_3^*(\alpha_{[27,2,1]}) - j_3^*(\alpha_{[21,8,2]}) + j_3^*(\alpha_{[3,2,9]}) + j_3^*(\alpha_{[9,2,3]}) - j_3^*(\alpha_{[6,-4,5]}) - j_3^*(\alpha_{[15,14,5]})) = -11968,$$

$$\frac{1}{\sqrt{21}}\mathbf{t}^{(3)}(21, 8) = \frac{1}{\sqrt{21}}(j_3^*(\alpha_{[42,0,1]}) - j_3^*(\alpha_{[21,0,2]}) + j_3^*(\alpha_{[3,0,14]}) - j_3^*(\alpha_{[6,0,7]})) = 342144.$$

We want to express these traces as the coefficients of $f_{d,p}$. As explained in the Appendix of [13], $f_{d,p}$ can be found by making use of “Rankin-Cohen bracket” $[,]_n$. Let $f(\tau)$ and $g(\tau)$ denote two modular forms of weight k and l on some group $\Gamma' \subset \Gamma$. We denote by D the differential operator $\frac{1}{2\pi i} \frac{d}{d\tau} = q \frac{d}{dq}$ and use $f', f'', \dots, f^{(n)}$ instead of $Df, D^2f, \dots, D^n f$. The n -th Rankin-Cohen bracket of f and g is defined by the formula

$$[f, g]_n(\tau) = \sum_{r+s=n} (-1)^r \binom{n+k-1}{s} \binom{n+l-1}{r} f^{(r)}(\tau) g^{(s)}(\tau).$$

By [8] §7 (or [17] §1) $[f, g]_n(\tau)$ is a modular form of weight $k+l+2n$ on Γ' . Let us consider the construction of $f_{d,p}$ in the cases $p = 2$ and $p = 3$:

$p = 2$: Let $\theta = \sum_{n \in \mathbb{Z}} q^{n^2}$, $u = ([\theta, E_{10}(8\tau)]_1 / \Delta(8\tau) + 2112\theta) / (-40)$ and $v = ([\theta, E_8(8\tau)]_2 / \Delta(8\tau) - 11520\theta) / 72$. Then it follows that

$$f_{4,2} = (v - u) / 12 = q^{-4} - 52q + 272q^4 + 2600q^8 - 8244q^9 + 15300q^{12} + 71552q^{16} - 204800q^{17} + 282880q^{20} + \dots,$$

$$f_{7,2} = (4u - v) / 3 = q^{-7} - 23q - 2048q^4 + 45056q^8 + 252q^9 - 516096q^{12} + 4145152q^{16} - 1771q^{17} - 26378240q^{20} + \dots$$

$p = 3$: Let $u = ([\theta, E_{10}(12\tau)]_1 / \Delta(12\tau) + 1584\theta) / (-20)$, $v = ([\theta, E_8(12\tau)]_2 / \Delta(12\tau) - 25920\theta) / 72$ and $w = ([\theta, E_6(12\tau)]_3 / \Delta(12\tau) + 272160\theta) / (-112)$. Then we obtain that

$$f_{3,3} = (4u - 5v + w) / 360 = q^{-3} - 14q + 40q^4 - 78q^9 + 168q^{12} - 378q^{13} + 688q^{16} - 897q^{21} \dots,$$

$$f_{8,3} = (-9u + 10v - w) / 60 = q^{-8} - 34q - 188q^4 + 2430q^9 + 8262q^{12} - 11968q^{13} - 34936q^{16} + 171072q^{21} \dots,$$

$$f_{11,3} = (36u - 13v + w) / 24 = q^{-11} + 22q - 552q^4 - 11178q^9 + 48600q^{12} + 76175q^{13} - 269744q^{16} - 1782891q^{21} \dots$$

Comparing these coefficients with the examples of twisted traces $\mathbf{t}^{(p)}(D, d)$ we are led to the following result which can be proved in a way analogous to the proof of [18] Theorem 6:

$$\frac{1}{\sqrt{D}} \mathbf{t}^{(p)}(D, d) = A^*(D, d) = -B^*(D, d) \quad (17)$$

with $A^*(D, d) = 2^{s(D,p)} A(D, d)$ and $B^*(D, d) = 2^{s(D,p)} B(D, d)$.

Now we will derive certain product identities from (17).

Lemma 4.2. *For $i \geq 0$ and m coprime to p ,*

$$\phi_{p^{2i}m^2D,1}|_{V_p} = p\phi_{p^{2i+2}m^2D,p} + \phi_{p^{2i}m^2D,p}.$$

Here V_p is the Hecke operator on Jacobi forms defined by the formula (2) in [11].

Proof. According to [11] Theorem 4.1, the operator V_p maps $J_{2,1}^1$ to $J_{2,p}^1$. From the formula (7) in [11], p.43, we find that

$$\text{the coefficient of } q^n \zeta^r \text{ in } \phi_{p^{2i}m^2D,1}|_{V_p} = \begin{cases} p, & \text{if } 4pn - r^2 = -p^{2i+2}m^2D \\ 1, & \text{if } 4pn - r^2 = -p^{2i}m^2D \\ 0, & \text{if } 4pn - r^2 < 0, \neq -p^{2i+2}m^2D, -p^{2i}m^2D. \end{cases}$$

From these observations and the uniqueness of $\phi_{D,p}$, the lemma immediately follows. $\square$

Let $t = j_p^*$. For each positive integer m , we define

$$\mathbf{t}_m^{(p)}(D, d) = \sum_{Q \in \mathcal{Q}_{dD,p,\beta}/\Gamma_0(p)} \chi(Q) t_m(\alpha_Q)$$

where t_m is a unique polynomial of t such that $t_m \equiv q^{-m} \pmod{q\mathbb{C}[[q]]}$. We can use Hecke operators in integral and half-integral weight to get the following lemma whose proof is similar to that of formulas in p.13 of [18].

Lemma 4.3. *Let l be a positive integer coprime to p and $d = 4pn - r^2$. Then*

- (i) $\frac{1}{\sqrt{D}} \mathbf{t}_l^{(p)}(D, d) = -\text{coefficient of } q^n \zeta^r \text{ in } 2^{s(D,p)} \phi_{D,p}|_{T_l},$
- (ii) $\phi_{D,p}|_{T_l} = \sum_{\nu|l} \left(\frac{D}{l/\nu} \right) \nu \phi_{\nu^2 D, p}.$

We claim that for $d = 4pn - r^2$,

$$\frac{1}{\sqrt{D}} \mathbf{t}_m^{(p)}(D, d) = -\text{coefficient of } q^n \zeta^r \text{ in } \sum_{u|m} 2^{s(u^2 D, p)} \left(\frac{D}{m/u} \right) u \phi_{u^2 D, p}. \quad (18)$$

Let $J = j - 744$ and J_m be the unique polynomial such that $J_m \equiv q^{-m} \pmod{q\mathbb{C}[[q]]}$. By “expansion formula” ([14] p.14 or [9] p.319) we come up with $t_{lp^{k+1}}(\tau) = J_{lp^k}(p\tau) + J_{lp^k}(\tau) - t_{lp^k}(\tau)$ (for $k \geq 0$) and therefore

$$\begin{aligned} & \sum_{Q \in \mathcal{Q}_{dD, p, \beta} / \Gamma_0(p)} \chi(Q) t_{lp^{k+1}}(\alpha_Q) \\ &= \sum_{Q \in \mathcal{Q}_{dD, p, \beta} / \Gamma_0(p)} \chi(Q) (J_{lp^k}(p\tau) + J_{lp^k}(\tau))|_{\tau=\alpha_Q} - \sum_{Q \in \mathcal{Q}_{dD, p, \beta} / \Gamma_0(p)} \chi(Q) t_{lp^k}(\alpha_Q). \end{aligned} \quad (19)$$

The map which sends $[a, b, c] \in \mathcal{Q}_{dD, p, \beta}$ to $[a/p, b, cp] \in \mathcal{Q}_{dD}$ induces a bijection between $\mathcal{Q}_{dD, p, \beta} / \Gamma_0(p)$ and $\mathcal{Q}_{dD} / \Gamma$. And the natural map from $\mathcal{Q}_{dD, p, \beta} / \Gamma_0(p)$ to $\mathcal{Q}_{dD} / \Gamma$ also gives a bijection. These maps preserve the values of χ and (19) is rewritten as

$$\sum_{Q \in \mathcal{Q}_{dD, p, \beta} / \Gamma_0(p)} \chi(Q) t_{lp^{k+1}}(\alpha_Q) = 2 \sum_{Q \in \mathcal{Q}_{dD} / \Gamma} \chi(Q) J_{lp^k}(\alpha_Q) - \sum_{Q \in \mathcal{Q}_{dD, p, \beta} / \Gamma_0(p)} \chi(Q) t_{lp^k}(\alpha_Q),$$

which yields

$$\mathbf{t}_{lp^{k+1}}^{(p)}(D, d) = 2\mathbf{t}_{lp^k}^{(1)}(D, d) - \mathbf{t}_{lp^k}^{(p)}(D, d) \quad \text{for } k \geq 0.$$

In (18) we write $m = lp^k$ with $(l, p) = 1$. We use induction on k to prove the claim. If $k = 0$, the claim (18) follows from Lemma 4.3. Now we assume the claim for k .

$$\begin{aligned}
\frac{1}{\sqrt{D}} \mathbf{t}_{lp^{k+1}}^{(p)}(D, d) &= \frac{1}{\sqrt{D}} (2\mathbf{t}_{lp^k}^{(1)}(D, d) - \mathbf{t}_{lp^k}^{(p)}(D, d)) \\
&= -\text{coefficient of } q^n \zeta^r \text{ in } 2(\phi_{D,1}|_{T_{lp^k}})|_{V_p} - \frac{1}{\sqrt{D}} \mathbf{t}_{lp^k}^{(p)}(D, d) \text{ by [18] formula (25), Theorem 5-(iii)} \\
&= -\text{coefficient of } q^n \zeta^r \text{ in } 2\left(\sum_{i=0}^k \sum_{\nu|l} \nu p^i \left(\frac{D}{(l/\nu)p^{k-i}}\right) \phi_{\nu^2 p^{2i} D, 1}\right)|_{V_p} - \frac{1}{\sqrt{D}} \mathbf{t}_{lp^k}^{(p)}(D, d) \\
&\quad \text{by [18] p.13 and Theorem 5-(iii)} \\
&= -\text{coefficient of } q^n \zeta^r \text{ in } \left[2 \sum_{i=0}^k \sum_{\nu|l} \nu p^i \left(\frac{D}{(l/\nu)p^{k-i}}\right) (p\phi_{\nu^2 p^{2i+2} D, p} + \phi_{\nu^2 p^{2i} D, p}) \right. \\
&\quad \left. - \left(\sum_{i=1}^k \sum_{\nu|l} 2\nu p^i \left(\frac{D}{(l/\nu)p^{k-i}}\right) \phi_{\nu^2 p^{2i} D, p} + \sum_{\nu|l} 2^{s(D,p)} \left(\frac{D}{(l/\nu)p^k}\right) \nu \phi_{\nu^2 D, p}\right) \right] \\
&\quad \text{by Lemma 4.2 and induction hypothesis} \\
&= -\text{coefficient of } q^n \zeta^r \text{ in } \left[\sum_{i=1}^{k+1} \sum_{\nu|l} 2\nu p^i \left(\frac{D}{(l/\nu)p^{k+1-i}}\right) \phi_{\nu^2 p^{2i} D, p} + \sum_{\nu|l} 2^{s(D,p)} \left(\frac{D}{(l/\nu)p^{k+1}}\right) \nu \phi_{\nu^2 D, p}\right] \\
&\quad \text{since } \left(\frac{D}{p}\right) = 0 \text{ or } 1, \text{ and } 2^{s(D,p)} = 1 \text{ or } 2.
\end{aligned}$$

Let $z \in \mathfrak{H}$. Note that $\frac{1}{m} t_m(z)$ can be viewed as the coefficient of q^m -term in $-\log q - \log(t(\tau) - t(z))$ (see [16]). Thus $\log q^{-1} - \sum_{m>0} \frac{1}{m} t_m(z) q^m = \log(t(\tau) - t(z))$. Taking exponential on both sides, we get

$$q^{-1} \exp\left(-\sum_{m>0} \frac{1}{m} t_m(z) q^m\right) = t(\tau) - t(z). \quad (20)$$

By the claim (18), we obtain

$$\frac{1}{\sqrt{D}} \mathbf{t}_m^{(p)}(D, d) = -\sum_{u|m} u \left(\frac{D}{m/u}\right) B^*(u^2 D, d). \quad (21)$$

From (20) and (21) it follows that

$$\begin{aligned}
 \prod_{Q \in \mathcal{Q}_{dD,p,\beta}/\Gamma_0(p)} (t(\tau) - t(\alpha_Q))^{\chi(Q)} &= \exp\left(-\sum_{m=1}^{\infty} \mathbf{t}_m^{(p)}(D, d) q^m/m\right) \\
 &= \exp\left(\sqrt{D} \sum_{m=1}^{\infty} \sum_{u|m} u \left(\frac{D}{m/u}\right) B^*(u^2 D, d) q^m/m\right) = \exp\left(\sqrt{D} \sum_{m=1}^{\infty} \sum_{u=1}^{\infty} u \left(\frac{D}{m}\right) B^*(u^2 D, d) \frac{q^{mu}}{mu}\right) \\
 &= \exp\left(\sum_{u=1}^{\infty} (-B^*(u^2 D, d)) \left(-\sum_{0 < n < D} \left(\frac{D}{n}\right) \sum_{m=1}^{\infty} (\zeta_D^n q^u)^m/m\right)\right) \\
 &\quad \text{by the Gauss sum identity } \sqrt{D} \left(\frac{D}{m}\right) = \sum_{0 < n < D} \left(\frac{D}{n}\right) \zeta_D^{nm} \\
 &= \exp\left(\sum_{u=1}^{\infty} A^*(u^2 D, d) \sum_{0 < n < D} \left(\frac{D}{n}\right) \log(1 - \zeta_D^n q^u)\right).
 \end{aligned}$$

Now the following theorem is immediate.

Theorem 4.4. *Let $D > 1$ and $-d < 0$ be coprime fundamental discriminants and congruent to a square modulo $4p$. We define $\mathcal{H}_{D,d,p}(X) = \prod_{Q \in \mathcal{Q}_{dD,p,\beta}/\Gamma_0(p)} (X - j_p^*(\alpha_Q))^{\chi(Q)}$. Then*

$$\mathcal{H}_{D,d,p}(j_p^*(\tau)) = \prod_{u=1}^{\infty} P_D(q^u)^{A^*(u^2 D, d)}$$

where $A^*(D, d) = 2^{s(D,p)} A(D, d)$ and $P_D(t) = \prod_{0 < n < D} (1 - \zeta_D^n t)^{\left(\frac{D}{n}\right)}$.

Example 4.5. Let us consider the case $p = 3$ and $D = 13, d = 3$. For $\beta \equiv 1 \pmod{6}$, a set of representatives for $\mathcal{Q}_{39,3,1}/\Gamma_0(3)$ is given by $\{[12, 3, 1], [3, 3, 4], [6, 3, 2], [15, 9, 2]\}$. Meanwhile the q -expansion of $f_{3,3}$ is

$$q^{-3} - 14q + 40q^4 - 78q^9 + 168q^{12} - 378q^{13} + 688q^{16} + \dots + 133056q^{52} + \dots + (-30650256)q^{117} + \dots$$

Theorem 4.4 yields the following product formula:

$$\frac{(j_3^*(\tau) - j_3^*(\alpha_{[12,3,1]})) (j_3^*(\tau) - j_3^*(\alpha_{[3,3,4]}))}{(j_3^*(\tau) - j_3^*(\alpha_{[6,3,2]})) (j_3^*(\tau) - j_3^*(\alpha_{[15,9,2]}))} = P_{13}(q)^{-378} P_{13}(q^2)^{133056} P_{13}(q^3)^{2(-30650256)} \dots$$

Acknowledgment

I would like to thank Professor J. H. Bruinier for encouraging me to do this subject and kindly answering my questions. I also want to express my gratitude to the people at Mathematisches Institut der Universität Heidelberg for their support.

REFERENCES

- [1] R. E. Borcherds, *Monstrous moonshine and monstrous Lie superalgebras*, Invent. math. **109** (1992), 405-444.
- [2] R. Borcherds, *Automorphic forms on $O_{s+2,2}(\mathbb{R})$ and infinite products*, Invent. Math. **120** (1995), 161-213.
- [3] R. Borcherds, *Automorphic forms with singularities on Grassmanians*, Invent. Math. **132** (1998), 491-562.
- [4] R. Borcherds, *The Gross-Kohnen-Zagier theorem in higher dimensions*, Duke Math. J. **97** (1999), 219-233.
- [5] J. H. Bruinier, *Borcherds products on $O(2, l)$ and Chern classes of Heegner divisors*, Springer Lecture Notes in Mathematics 1780, Springer-Verlag (2002).
- [6] J. H. Bruinier and M. Bundschuh, *On Borcherds products associated with lattices of prime discriminant* to appear in Ramanujan J.
- [7] I. Chen and N. Yui, *Singular values of Thompson series*, in “Groups, Difference Sets and Monster” (K. T. Arasu *et al.* Eds.), pp. 255-326, de Gruyter, Berlin, 1995.
- [8] H. Cohen, *Sums involving the values at negative integers of L -functions of quadratic characters*, Math. Ann. **217** (1975), 271-285.
- [9] J. H. Conway and S. P. Norton, *Monstrous Moonshine*, Bull. London Math. Soc. **11**, 308-339, 1979.
- [10] D. Cox, *Primes of the form $x^2 + ny^2$* , John Wiley & Sons, 1989.
- [11] M. Eichler and D. Zagier, *The Theory of Jacobi Forms*, Progress in Math. **55**, Birkhäuser-Verlag, Boston-Basel-Stuttgart, 1985.
- [12] B. Gross, W. Kohnen, D. Zagier, *Heegner points and derivatives of L -series II*, Math. Ann. **278** (1987), 497-562.
- [13] C. H. Kim, *Borcherds products associated with certain Thompson series*, to appear in Compositio Math., e-print arXiv:math.NT/0203076.
- [14] M. Koike, *On replication formula and Hecke operators*, Nagoya University (preprint).

- [15] S. Lang, *Algebraic number theory*, (2nd ed.), Springer-Verlag, 1994.
- [16] S. P. Norton, *More on moonshine*. In: Computational Group Theory, 185-193, Academic Press, 1984.
- [17] D. Zagier, *Modular forms and differential operators*, Proc. Indian Acad. Sci. **104** (1994), 57-75.
- [18] D. Zagier, *Traces of singular moduli*, Max-Planck-Institut für Mathematik, Preprint series **2000** (8).

MATHEMATISCHES INSTITUT DER UNIVERSITÄT HEIDELBERG, IM NEUENHEIMER FELD 288, 69120
HEIDELBERG

E-mail address: kim@mathi.uni-heidelberg.de